

Homework 4

Problem 1. (a) Prove the “Squeeze Rule”: If $0 \leq X_n \leq Y_n$ and $Y_n \rightarrow_p 0$, then $X_n \rightarrow_p 0$; (b) Prove: $X_n \rightarrow_p 0$ if and only if $|X_n| \rightarrow_p 0$.

Solution. (a) For any $\epsilon > 0$,

$$\Pr(Y_n \leq \epsilon) \leq \Pr(X_n \leq \epsilon) \leq 1.$$

Then,

$$\Pr(Y_n \leq \epsilon) = \Pr(|Y_n - 0| \leq \epsilon) \rightarrow 1 \implies \Pr(X_n \leq \epsilon) = \Pr(|X_n - 0| \leq \epsilon) \rightarrow 1.$$

(b) “ \implies part”: by continuous mapping theorem, since the mapping $x \mapsto |x|$ is continuous. “ \impliedby part”: straightforward.

Problem 2. Provide a counter example to show that $X_n \rightarrow_d X$ and $Y_n \rightarrow_d Y$ does not imply $X_n + Y_n \rightarrow_d X + Y$. Hint: Consider an iid random sample X_1, \dots, X_n with $\mathbb{E}X_1 = 0$ and $n^{1/2}\bar{X}_n$ and $-n^{1/2}\bar{X}_n$.

Solution. Let Z be a random variable such that $Z \sim N(0, \sigma^2)$, where $\sigma^2 = \text{Var}(X_1)$. Then by CLT,

$$n^{1/2}\bar{X}_n \rightarrow_d Z$$

and

$$-n^{1/2}\bar{X}_n = (-1) \times (n^{1/2}\bar{X}_n) \rightarrow_d (-1) \times Z \sim N(0, \sigma^2).$$

Therefore, it is also true that $-n^{1/2}\bar{X}_n \rightarrow_d Z \sim N(0, \sigma^2)$. Note

$$0 = (n^{1/2}\bar{X}_n) + (-n^{1/2}\bar{X}_n) \rightarrow_d Z + Z \sim N(0, 4\sigma^2).$$

Problem 3. Let $\hat{\theta}_n = (\hat{\theta}_{n,1}, \dots, \hat{\theta}_{n,k})'$ be an estimator of the k -vector of parameters $\theta = (\theta_1, \dots, \theta_k)'$.

Suppose that $\hat{\theta}_n \rightarrow_p \theta$, and $n^{1/2}(\hat{\theta}_n - \theta) \rightarrow_d W \sim N(0, \Sigma)$, where Σ is a positive definite $k \times k$ matrix. Use the delta method or CMT to find the (non-degenerate, i.e., not a constant) asymptotic distributions of the following quantities after a suitable normalization. "Suitable normalization" means subtraction of a constant and/or multiplication by a constant (could be dependent on n).

1. $n^{1/2}(\hat{\theta}_n - \theta)' \mathbf{c}$ where $\mathbf{c} \in \mathbb{R}^k$ is a vector of constants.
2. $\hat{\theta}_{n,1}$.
3. $n(\hat{\theta}_n - \theta)'(\hat{\theta}_n - \theta)$.
4. $\hat{\theta}_{n,1} - \hat{\theta}_{n,2}$.
5. $\hat{\theta}_{n,1}\hat{\theta}_{n,2}/\hat{\theta}_{n,3}$, provided that $\theta_3 \neq 0$.

Solution.

1. Define $\mathbf{X}_n = n^{1/2}(\hat{\theta}_n - \theta)$ and $h(\mathbf{X}_n) = \mathbf{X}_n' \mathbf{c}$. By the Continuous Mapping Theorem we have

$$n^{1/2}(\hat{\theta}_n - \theta)' \mathbf{c} = h(\mathbf{X}_n) \rightarrow_d h(\mathbf{W}) = \mathbf{W}' \mathbf{c}$$

By the property of normal distribution we have

$$n^{1/2}(\hat{\theta}_n - \theta)' \mathbf{c} \rightarrow_d \mathbf{W}' \mathbf{c} \sim N(0, \mathbf{c}' \Sigma \mathbf{c}).$$

2. Set $\mathbf{c} = (1, 0, \dots, 0)'$. Then, it follows from Part (i) that

$$n^{1/2}(\hat{\theta}_{n,1} - \theta_1) \rightarrow_d N(0, \sigma_{11}^2),$$

where σ_{11}^2 is the first diagonal element of Σ .

3. Since $n^{1/2}(\hat{\theta}_n - \theta) \rightarrow_d \mathbf{W}$, by the Continuous Mapping Theorem,

$$n(\hat{\theta}_n - \theta)'(\hat{\theta}_n - \theta) = \left[n^{1/2}(\hat{\theta}_n - \theta) \right]' \left[n^{1/2}(\hat{\theta}_n - \theta) \right] \rightarrow_d \mathbf{W}'\mathbf{W}.$$

4. Set $\mathbf{c} = (1, -1, 0, \dots, 0)'$. It follows from Part (i) that

$$n^{1/2}(\hat{\theta}_{n,1} - \hat{\theta}_{n,2} - \theta_1 + \theta_2) \rightarrow_d N(0, \sigma_{11}^2 - 2\sigma_{12} + \sigma_{22}^2),$$

where σ_{11}^2 and σ_{22}^2 are the first and second diagonal element of Σ , and σ_{12} is the element on the first row and second column of Σ .

5. Put $h(\theta) = \frac{\theta_1\theta_2}{\theta_3}$, apply the Delta method

$$n^{1/2}\left(\frac{\hat{\theta}_{n,1}\hat{\theta}_{n,2}}{\hat{\theta}_{n,3}} - \frac{\theta_1\theta_2}{\theta_3}\right) = n^{1/2}(h(\hat{\theta}_n) - h(\theta)) \rightarrow_d \frac{\partial h(\theta)}{\partial \theta'} \mathbf{W}$$

where

$$\frac{\partial h(\theta)}{\partial \theta'} = \left(\frac{\theta_2}{\theta_3}, \frac{\theta_1}{\theta_3}, \frac{-\theta_1\theta_2}{\theta_3^2}, 0, \dots, 0 \right)'.$$

Then by the property of Normal density

$$n^{1/2}\left(\frac{\hat{\theta}_{n,1}\hat{\theta}_{n,2}}{\hat{\theta}_{n,3}} - \frac{\theta_1\theta_2}{\theta_3}\right) \rightarrow_d N\left(0, \frac{\partial h(\theta)}{\partial \theta'} \Sigma \frac{\partial h(\theta)'}{\partial \theta}\right).$$

Problem 4. Suppose that $\hat{\theta}_n \rightarrow_p \theta$ and $\hat{\beta}_n \rightarrow \beta$, where θ and β are two scalar parameters. Without relying on Slutsky's Theorem, show:

1. $c\hat{\theta}_n \rightarrow_p c\theta$, where c is a constant.

2. $\hat{\theta}_n\hat{\beta}_n \rightarrow_p \theta\beta$.

Solution. (i) Suppose $c \neq 0$. Then $\Pr(|c\hat{\theta}_n - c\theta| > \varepsilon) = \Pr(|\hat{\theta}_n - \theta| > \frac{\varepsilon}{|c|}) \rightarrow 0$ as $n \rightarrow \infty$. If $c = 0$, then $c\hat{\theta}_n = 0 \rightarrow_p c\theta = 0$.

(ii) First, note that $\hat{\theta}_n\hat{\beta}_n - \theta\beta = (\hat{\theta}_n - \theta + \theta)(\hat{\beta}_n - \beta + \beta) - \theta\beta = (\hat{\theta}_n - \theta)(\hat{\beta}_n - \beta) + (\hat{\theta}_n - \theta)\beta + (\hat{\beta}_n - \beta)\theta$. Then, $(\hat{\theta}_n - \theta)\beta + (\hat{\beta}_n - \beta)\theta \rightarrow_p 0$ by Part (i). Then, for any $\epsilon > 0$,

$$\begin{aligned} \Pr\left(|(\hat{\theta}_n - \theta)(\hat{\beta}_n - \beta)| > \varepsilon\right) &\leq \Pr\left(|\hat{\theta}_n - \theta| > \sqrt{\varepsilon} \text{ or } |\hat{\beta}_n - \beta| > \sqrt{\varepsilon}\right) \\ &\leq \Pr\left(|\hat{\theta}_n - \theta| > \sqrt{\varepsilon}\right) + \Pr\left(|\hat{\beta}_n - \beta| > \sqrt{\varepsilon}\right) \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Thus, $\hat{\theta}_n\hat{\beta}_n - \theta\beta \rightarrow_p 0$.

Problem 5. Suppose that $\mathbb{E}(\hat{\theta}_n) \rightarrow \theta$ and $\text{Var}(\hat{\theta}_n) \rightarrow 0$ as $n \rightarrow \infty$. Show that $\hat{\theta}_n \rightarrow_p \theta$.

Solution. $\hat{\theta}_n$ converges in probability to θ if for all $\varepsilon > 0$, $\Pr\left(\left|\hat{\theta}_n - \theta\right| \geq \varepsilon\right) \rightarrow 0$ as $n \rightarrow \infty$. First, decompose the Mean Squared Error (MSE) into

$$\begin{aligned} MSE\left(\hat{\theta}_n\right) &= \mathbb{E}\left(\hat{\theta}_n - \theta\right)^2 = \mathbb{E}\left(\hat{\theta}_n - \mathbb{E}\hat{\theta}_n + \mathbb{E}\hat{\theta}_n - \theta\right)^2 \\ &= \mathbb{E}\left(\hat{\theta}_n - \mathbb{E}\hat{\theta}_n\right)^2 + \left(\mathbb{E}\hat{\theta}_n - \theta\right)^2 + 2\mathbb{E}\left(\hat{\theta}_n - \mathbb{E}\hat{\theta}_n\right)\left(\mathbb{E}\hat{\theta}_n - \theta\right) \\ &= \mathbb{E}\left(\hat{\theta}_n - \mathbb{E}\hat{\theta}_n\right)^2 + \left(\mathbb{E}\hat{\theta}_n - \theta\right)^2 = \text{Var}\left(\hat{\theta}_n\right) + \text{Bias}\left(\hat{\theta}_n\right)^2, \end{aligned}$$

where the last line follows by the fact that $\mathbb{E}\left(\hat{\theta}_n - \mathbb{E}\hat{\theta}_n\right) = 0$.

Then, using Markov's Inequality,

$$\Pr\left(\left|\hat{\theta}_n - \theta\right| \geq \varepsilon\right) \leq \frac{\mathbb{E}\left|\hat{\theta}_n - \theta\right|^2}{\varepsilon^2} = \frac{\mathbb{E}\left(\hat{\theta}_n - \theta\right)^2}{\varepsilon^2} = \frac{\text{Var}\left(\hat{\theta}_n\right) + \text{Bias}\left(\hat{\theta}_n\right)^2}{\varepsilon^2} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

since by assumption, $\text{Var}\left(\hat{\theta}_n\right) \rightarrow 0$ and $\mathbb{E}\hat{\theta}_n - \theta \rightarrow 0$ as $n \rightarrow \infty$.

Problem 6. Consider the linear model (with independently and identically distributed (i.i.d.) observations):

$$Y_i = \beta_0 + \beta_1 X_{1,i} + \beta_2 X_{2,i} + U_i$$

with $\mathbb{E}U_i = \mathbb{E}U_i X_{1,i} = \mathbb{E}U_i X_{2,i} = 0$. Suppose we know that $\beta_2 = \beta_1$ and conduct a constrained LS estimation of β_1 :

$$\min_{b_0, b_1} \sum_{i=1}^n (Y_i - b_0 - b_1 X_{1,i} - b_1 X_{2,i})^2.$$

1. Find the expression for the constrained LS estimator $(\hat{\beta}_0, \hat{\beta}_1)$ that solve the above minimization problem.
2. Assume that the restriction $\beta_2 = \beta_1$ is true. Derive the large-sample (asymptotic) distribution of $\hat{\beta}_1$.

Solution. Denote $\bar{X}_1 = n^{-1} \sum_{i=1}^n X_{1,i}$ and $\bar{X}_2 = n^{-1} \sum_{i=1}^n X_{2,i}$. The constrained LS:

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2) Y_i}{\sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2)^2}.$$

And

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 (\bar{X}_1 + \bar{X}_2).$$

For (ii),

$$\begin{aligned} \hat{\beta}_1 &= \frac{\sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2) Y_i}{\sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2)^2} \\ &= \frac{\sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2) (\beta_0 + \beta_1 X_{1,i} + \beta_1 X_{2,i} + U_i)}{\sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2)^2} \\ &= \beta_1 + \frac{\frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2) U_i}{\frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2)^2}. \end{aligned}$$

By WLLN and Continuous Mapping Theorem,

$$\frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2)^2 = \frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i})^2 - (\bar{X}_1 + \bar{X}_2)^2$$

$$\rightarrow_p \text{Var}(X_{1,i} + X_{2,i}).$$

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2) U_i &= \frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \mathbb{E}(X_{1,i} + X_{2,i})) U_i \\ &\quad + (\bar{X}_1 - \mathbb{E}(X_{1,i})) \frac{1}{n} \sum_{i=1}^n U_i + (\bar{X}_2 - \mathbb{E}(X_{2,i})) \frac{1}{n} \sum_{i=1}^n U_i. \end{aligned}$$

Since $n^{-1/2} \sum_{i=1}^n U_i \rightarrow_d N(0, \mathbb{E}(U_i^2))$, $\bar{X}_1 - \mathbb{E}(X_{1,i}) \rightarrow_p 0$ and $\bar{X}_2 - \mathbb{E}(X_{2,i}) \rightarrow_p 0$, by Slutsky's theorem,

$$\begin{aligned} (\bar{X}_1 - \mathbb{E}(X_{1,i})) \frac{1}{\sqrt{n}} \sum_{i=1}^n U_i &\rightarrow_p 0 \\ (\bar{X}_2 - \mathbb{E}(X_{2,i})) \frac{1}{\sqrt{n}} \sum_{i=1}^n U_i &\rightarrow_p 0. \end{aligned}$$

By CLT,

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \mathbb{E}(X_{1,i} + X_{2,i})) U_i \rightarrow_d N\left(0, \mathbb{E}\left(U_i^2 (X_{1,i} + X_{2,i} - \mathbb{E}(X_{1,i} + X_{2,i}))^2\right)\right).$$

By Slutsky's theorem,

$$\begin{aligned} \sqrt{n}(\hat{\beta}_1 - \beta_1) &= \frac{\frac{1}{\sqrt{n}} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2) U_i}{\frac{1}{n} \sum_{i=1}^n (X_{1,i} + X_{2,i} - \bar{X}_1 - \bar{X}_2)^2} \\ &\rightarrow_d \text{Var}(X_{1,i} + X_{2,i})^{-1} N\left(0, \mathbb{E}\left(U_i^2 (X_{1,i} + X_{2,i} - \mathbb{E}(X_{1,i} + X_{2,i}))^2\right)\right). \end{aligned}$$

Problem 7. Suppose we observe the i.i.d. random sample $\{(Y_i, X_i)\}_{i=1}^n$ with X_i being a scalar. Take the linear model

$$\begin{aligned} Y_i &= X_i \beta + e_i \\ \mathbb{E}(e_i | X_i) &= 0. \end{aligned}$$

Consider the estimator

$$\hat{\beta} = \frac{\sum_{i=1}^n X_i^3 Y_i}{\sum_{i=1}^n X_i^4}.$$

Find the asymptotic distribution of $\sqrt{n}(\hat{\beta} - \beta)$.

Problem 8. Let $\{\theta_n : n \geq 1\}$ be a random sequence such that $\Pr(\theta_n = 0) = (n-1)/n$, and $\Pr(\theta_n = n^2) = 1/n$. Note that the only possible values for θ_n are zero and n^2 .

1. Show that $\lim_{n \rightarrow \infty} \mathbb{E}\theta_n = \infty$.
2. Does θ_n converge in probability to some limit? If yes, prove. If not, explain why.

Solution. (i) $\mathbb{E}\theta_n = 0 \cdot (n-1)/n + n^2 \cdot 1/n = n \rightarrow \infty$.

(ii) $\theta_n \rightarrow_p 0$, since for any $\epsilon > 0$, $\Pr(|\theta_n| > \epsilon) = \Pr(\theta_n = n^2) = 1/n \rightarrow 0$.